

**2019-novel coronavirus (2019-nCoV) infections trigger an exaggerated cytokine response aggravating lung injury**

Yingxia Liu<sup>1,5</sup>, Cong Zhang<sup>2,3,5</sup>, Fengming Huang<sup>3,5</sup>, Yang Yang<sup>1,5</sup>, Fuxiang Wang<sup>1</sup>, Jing Yuan<sup>1</sup>, Zheng Zhang<sup>1</sup>, Yuhao Qin<sup>3</sup>, Xiaoyun Li<sup>3</sup>, Dandan Zhao<sup>3</sup>, Shunwang Li<sup>3</sup>, Shuguang Tan<sup>4</sup>, Zhaoqin Wang<sup>1</sup>, Jinxiu Li<sup>1</sup>, Chenguang Shen<sup>1</sup>, Jianming Li<sup>1</sup>, Ling Peng<sup>1</sup>, Weibo Wu<sup>1</sup>, Mengli Cao<sup>1</sup>, Li Xing<sup>1</sup>, Zhixiang Xu<sup>1</sup>, Li Chen<sup>1</sup>, Congzhao Zhou<sup>2</sup>, William J. Liu<sup>4\*</sup>, Lei Liu<sup>1\*</sup>, Chengyu Jiang<sup>3\*</sup>

<sup>1</sup>Shenzhen Key Laboratory of Pathogen and Immunity, National Clinical Research Center for infectious disease, State Key Discipline of Infectious Disease, Shenzhen Third People's Hospital, Second Hospital Affiliated to Southern University of Science and Technology, Shenzhen, China.

<sup>2</sup>Hefei National Laboratory for Physical Sciences at the Microscale and School of Life Sciences, University of Science and Technology of China, Hefei, China.

<sup>3</sup>The State Key Laboratory of Medical Molecular Biology, Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences, Department of Biochemistry, Peking Union Medical College, Beijing, China.

<sup>4</sup>The NHC Key Laboratory of Biosafety, National Institute for Viral Disease Control and Prevention, China CDC, Beijing, China

<sup>5</sup> These authors contributed equally.

\*email: [Jiang@pumc.edu.cn](mailto:Jiang@pumc.edu.cn); [liulei3322@aliyun.com](mailto:liulei3322@aliyun.com); [liujun@ivdc.chinacdc.cn](mailto:liujun@ivdc.chinacdc.cn)

A recent outbreak of pneumonia in Wuhan, China was caused by 2019 novel coronavirus (2019-nCoV). Here we reported 12 patients with 2019-nCoV infections in Shenzhen, China; all of them developed pneumonia and half developed acute respiratory distress syndrome (ARDS). We demonstrated the plasma cytokine profiles of these 12 patients. Thirty-eight out of 48 cytokines measured in the plasmas of 2019-nCoV infected patients were significantly elevated compared to healthy individuals. Seventeen cytokines were linked to 2019-nCoV load. Fifteen cytokines, M-CSF, IL-10, IFN- $\alpha$ 2, IL-17, IL-4, IP-10, IL-7, IL-1ra, G-CSF, IL-12, IFN- $\gamma$ , IL-1 $\alpha$ , IL-2, HGF, and PDGF-BB, were strongly associated with lung injury Murray score, and could predict disease severity of 2019-nCoV infections according to area under the curve (AUC) of the receiver-operating characteristics (ROC) calculations. Our results suggest that 2019-nCoV infections trigger extensive changes in a wide array of cytokines, some of these cytokines could be potential biomarkers of disease severity of 2019-nCoV infections. These findings improve our understanding of the immunopathologic mechanisms of this emerging and still evolving disease and suggest that modulators of cytokine responses could play a therapeutic role in combating the disease.

Since the outbreak of 2019 novel coronavirus (2019-nCoV) infections in late December, 2019 in Wuhan, Hubei Province, China<sup>1-9</sup>, the number of infected persons has exceeded 10,000 and already surpassed that of the previous severe acute respiratory syndrome (SARS) epidemic<sup>10</sup>. Hitherto, reflecting the geographical reach of the pandemic-prone coronaviruses, 2019-nCoV-infected cases have been reported in Thailand<sup>11</sup>, Germany<sup>12</sup>, the USA<sup>13</sup> and other parts of the world. The mortality rate is 15% for a case series of 41 hospitalized patients with 2019-nCoV infections in Wuhan, China<sup>3</sup>, compared with 10% for SARS-CoV and 37% for Middle East respiratory syndrome coronavirus (MERS-CoV)<sup>14</sup>. Although 2019-nCoV, the seventh reported human-infecting member of the family Coronaviridae, which also includes SARS-CoV<sup>15</sup> and MERS-CoV<sup>16,17</sup>, has been identified as the causative agent, the immunopathologic mechanisms of 2019-nCoV-associated diseases have not been elucidated.

One prominent feature of 2019-nCoV infections is the presence of acute respiratory distress syndrome (ARDS) in a sizable proportion of the documented 2019-nCoV-infected cases: 29% (12/41) in the series by Huang *et al*<sup>3</sup>. and 17% in a series of 99 cases of 2019-nCoV pneumonia<sup>18</sup>. An elevated production of proinflammatory cytokines/chemokines<sup>19</sup> or even hypercytokinemia<sup>20,21</sup>, also known as cytokine storm, is present in SARS-CoV and MERS-CoV infections<sup>15,16,22</sup> and contributes to acute lung injury and development of ARDS<sup>19</sup>. In this study, we examined the plasma cytokine/chemokine profile of 12 patients with laboratory-confirmed 2019-nCoV infections in Shenzhen, China *versus* 8 healthy subjects, 8 bacterial pneumonia patients, and 8 patients with influenza virus A H7N9 infections (see Extended Data Table 1).

Analysis using the Bio-Plex Pro Human Cytokine Screening Panel revealed that extensive and significant elevations in 38 out of 48 plasma cytokines/chemokines in patients with severe 2019-nCoV pneumonia (2019-nCoV-S) *versus* healthy individuals, suggesting the occurrence of hypercytokinemia in 2019-nCoV-S patients (see Extended Data Table 2). Overall, the plasma cytokine/chemokine levels in 2019-nCoV-S patients were markedly lower than H7N9 influenza virus A-infected patients, but moderately higher than patients with bacterial pneumonia (see Extended Data Table 2). In addition, the plasma cytokine/chemokine levels were higher at week 2 than week 1 from symptomatic onset of 2019-nCoV infections (see Extended Data Table 3). The plasma cytokine/chemokine levels were comparable between days 8-14 since illness onset and day 15 after illness onset, indicating persistent elevations of plasma cytokines/chemokines at the later stage of the disease (see Extended Data Table 3), which are consistent with clinical features of 2019-nCoV infections <sup>1,2,23</sup>.

We quantified viral RNA loads in throat swabs, sputum, and bronchoalveolar lavage fluid (BALF) samples by quantitative reverse transcription polymerase chain reaction (qRT-PCR) (see Extended Data Table 4). Our Spearman correlation analysis revealed that 2019-nCoV viral load was highly positively associated with the plasma levels of 16 cytokines (M-CSF, IL-10, IFN- $\alpha$ 2, IL-13, IL-17, IL-4, IP-10, IL-1 $\beta$ , IL-7, IL-1ra, G-CSF, IL-12, IFN- $\gamma$ , IL-1 $\alpha$ , IL-2, and HGF), and negatively associated with PDGF-BB (Table 1). The findings suggest that 2019-nCoV infection was associated with an elevated production of a wide array of cytokines/chemokines in the plasma of 2019-nCoV infected patients.

Using Spearman rank coefficient correlation analysis, we discovered a strong positive linear association between the plasma levels of 15 cytokines (IL-12, IFN- $\gamma$ , IL-2, HGF, IFN- $\alpha$ 2, IL-4, IL-17, IP-10, G-CSF, IL-10, IL-1ra, M-CSF, IL-1 $\alpha$ , and IL-7) in 2019-nCoV-infected patients and lung injury Murray score and a negative association between PDGF-BB and Murray score (Fig. 1). The area under the curve (AUC) of the receiver operating characteristic (ROC) was above 0.8 for each of these 15 cytokines, indicating that these cytokines could predict disease severity of 2019-nCoV infections (Fig. 2). Furthermore, the plasma cytokines/chemokines levels in 2019-nCoV-S patients were significantly different than 2019-nCoV-M patients (Fig. 3). Our data suggested that these 15 cytokines may be biomarkers for disease severity in 2019-nCoV infected patients.

In summary, we have demonstrated that hypercytokinemia occurred in 2019-nCoV infected patients. We have discovered that 15 cytokines are linearly associated with lung injury (Murray score) and may be potential biomarkers for disease severity. These cytokines include anti-viral cytokines IFN- $\alpha$ 2 and IFN- $\gamma$ , IL1ra, IL2, 4, 7, 10, 12 and 17, chemokine IP-10, as well as G-CSF and M-CSF. It is interesting that the levels of proinflammatory cytokines Th1, Th2, and Th17 were all increased. Previous studies have demonstrated marked elevation of IP-10, IL-6, IL-8, MCP1, and MIP-1 $\alpha$  in serums from SARS-CoV infected patients<sup>15,24</sup>, and a significant increase of IL-10, 15, and 17 as well as IFN  $\alpha$ 2 and  $\gamma$  in plasma samples collected from MERS-CoV infected patients<sup>16,17</sup>. Although measurements of plasma cytokines were not performed at the same time, the increased level of cytokines in blood samples from SARS-CoV<sup>15,24</sup> and

MERS-CoV<sup>17</sup> infected patients appeared considerably higher than levels in 2019-nCoV infected patients. The most extreme case of hypercytokinemia detected in humans may come from serum samples of patients infected with avian influenza A virus. In some cases, hypercytokinemia factors have been associated with; IP-10, MCP-1, MIG, and IL-8 of patients infected with H5N1 virus<sup>25,26</sup>, as well as MIF, SCF, MCP-1, HGF, SCGF-beta, IP-10, IL-18, and IFN- $\gamma$  in patients infected with H7N9 virus<sup>24,27</sup>.

Although the mechanisms of cytokine-mediated communication are largely unknown, attempts to use cytokines or cytokine inhibitors therapeutically have been increasingly successful<sup>28,29</sup>. Interferon  $\alpha$  and  $\gamma$  have been frequently used clinically despite their unpleasant side effects<sup>30-32</sup>, and interferon  $\gamma$  is an officially recommended drug for treatment of 2019-nCoV infected patients in China National Health Commission Guidelines for Diagnosis and Treatment of 2019-nCoV Pneumonia<sup>33</sup>. A number of antagonistic antibodies to cytokines have been used clinically or in clinical trials for the treatment of autoimmune or autoinflammatory diseases (including monoclonal antibodies against IL-1<sup>34</sup>, IL-10<sup>28</sup>, IL-12<sup>35</sup>, IL-17<sup>36</sup>, and IP-10<sup>37</sup>). These antibodies can be repurposed to attenuate hypercytokinemia in 2019-nCoV infected patients and may provide potential treatment for the current outbreak of 2019-CoV.

## **METHODS**

### **Experimental ethics policy**

The study protocol was approved by the Ethics Committees of Shenzhen Third People's

Hospital (SZTHEC2016001). Verbal informed consents were obtained from all patients or patients family members. The study was conducted in accordance with the International Conference on Harmonisation Guidelines for Good Clinical Practice and the Declaration of Helsinki and institutional ethics guidelines.

### **Acquisition of clinical specimens**

Throat swabs, rectal swabs, and sputum, and BALF specimens were collected within 24 hours of blood sample collection from laboratory-confirmed 2019-nCoV cases upon admission in January 2020 and at various time points thereafter. Plasma samples were collected from 8 healthy subjects undergoing wellness examination in the hospital in the interim. In addition, we obtained archived plasma samples from 8 patients with laboratory-confirmed H7N9 infections who were hospitalized between January 2015 and March 2017, and 8 bacterial pneumonia patients who were hospitalized between August and December 2019.

### **qRT-PCR**

Viral RNAs were extracted from clinical specimens using the QIAamp RNA Viral Kit (Qiagen, Heiden, Germany) as instructed by the manufacturer. They were amplified by quantitative reverse transcription polymerase chain reaction (qRT-PCR) using primers and probes recommended by Chinese Center for Diseases Control and Prevention (China CDC)<sup>38</sup>, and a commercially available kit for 2019-nCoV detection (GeneoDX Co., Shanghai, China). Samples with a cycle threshold ( $C_t$ ) value  $\leq 38.0$  were

considered putatively positive. Samples whose  $C_t$  was higher than 38 were re-tested and considered positive if  $C_t$  was  $\geq 38$  but  $\leq 40$  and negative if viral RNAs were undetectable on the second test.

### **Disease severity classification and Murray Score calculation**

Severity of 2019-nCoV infection was graded according to China National Health Commission Guidelines for Diagnosis and Treatment of 2019-nCoV infection. Briefly, a patient was considered to have mild 2019-nCoV pneumonia (2019-nCoV-M) if he or she had fever, respiratory manifestations and radiological findings indicative of pneumonia. A patient was considered to have severe 2019-nCoV pneumonia (2019-nCoV-S) if he or she met any of the following: 1) respiratory distraction (respiration rate  $\geq 30$ /min; 2) resting oxygen saturation  $\leq 93\%$ , or 3) arterial oxygen partial pressure ( $PaO_2$ )/ fraction of inspired oxygen ( $FiO_2$ )  $\leq 300$  mmHg (1 mmHg = 0.133 kPa). a patient was considered to have critical 2019-nCoV pneumonia (2019-nCoV-C) if he or she had any of the following: 1) respiratory failure requiring mechanical ventilation, 2) shock, or 3) failure of other organs requiring ICU care. Clinicopathological variables of 12 patients with 2019-nCoV infections including 4 2019-nCoV-M patients, 5 2019-nCoV-S patients and 3 2019-nCoV-C patients were collected at admission, and disease severity was assessed using Murray scores<sup>39</sup>.

### **Cytokine and chemokine measurements**

Plasma concentrations of 48 cytokines and chemokines were measured in duplicate



using Bio-Plex Pro Human Cytokine Screening Panel (48-Plex #12007283, Bio-Rad) according to the manufacturer's instructions. The plasma samples were fixed in 2% paraformaldehyde before analysis and measured in biosafety level III laboratory.

### Statistical analysis

The Spearman rank correlation coefficient was used for linear correlation analysis between plasma cytokine levels and Murray score of patients with 2019-nCoV infections. The area under the receiver operating characteristic (ROC) curve (AUC) of plasma cytokine levels was estimated for 2019-nCoV-M and 2019-nCoV-S infections. ANOVA or Mann–Whitney U test were used to compare plasma cytokine levels among the groups. All statistical tests were calculated using SPSS 16.0 for Windows (SPSS, Inc., Chicago, IL, USA). A two-tailed P value of less than 0.05 was considered to be statistically significant.

### References

1. Zhu, N., *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *The New England journal of medicine* (2020).
2. Chan, J.F.-W., *et al.* A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *The Lancet* (2020).
3. Huang, C., *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* (2020).
4. Huang, C., *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* (2020).
5. Lu, H., Stratton, C.W. & Tang, Y.W. Outbreak of Pneumonia of Unknown Etiology in Wuhan China: the Mystery and the Miracle. *J Med Virol* (2020).
6. Tan WJ, Z.X., Ma XJ, *et al.* A novel coronavirus genome identified in a cluster of pneumonia cases — Wuhan, China 2019–2020. *China CDC Weekly* (2020).
7. Xintian Xu, P.C., Jingfang Wang, Jiannan Feng, Hui Zhou, Xuan Li, Wu Zhong and Pei Hao. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of

- its spike protein for risk of human transmission. *Science China Life Sciences* (2020).
8. Zhang, N., *et al.* Recent advances in the detection of respiratory virus infection in humans. *J Med Virol* (2020).
9. Hui, D.S., *et al.* The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health - The latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis* **91**, 264-266 (2020).
10. <http://weekly.chinacdc.cn/news/TrackingtheEpidemic.htm#NHCFeb01> (accessed Feb 1, 2020).
11. Sookaromdee P., W.V. Imported cases of 2019-novel coronavirus (2019-nCoV) in Thailand: Mathematical modeling of the outbreak. . *Asian Pac J Trop Med.* (2020.).
12. Rothe, C., *et al.* Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *N Engl J Med* (2020).
13. Holshue, M.L., *et al.* First Case of 2019 Novel Coronavirus in the United States. *N Engl J Med* (2020).
14. Zaki, A.M., van Boheemen, S., Bestebroer, T.M., Osterhaus, A.D. & Fouchier, R.A. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**, 1814-1820 (2012).
15. Jiang, Y., *et al.* Characterization of cytokine/chemokine profiles of severe acute respiratory syndrome. *Am J Respir Crit Care Med* **171**, 850-857 (2005).
16. Mahallawi, W.H., Khabour, O.F., Zhang, Q., Makhdom, H.M. & Suliman, B.A. MERS-CoV infection in humans is associated with a pro-inflammatory Th1 and Th17 cytokine profile. *Cytokine* **104**, 8-13 (2018).
17. Niu, P., *et al.* A novel human mAb (MERS-GD27) provides prophylactic and postexposure efficacy in MERS-CoV susceptible mice. *Sci China Life Sci* **61**, 1280-1282 (2018).
18. Chen, N., *et al.* Epidemiological and Clinical Characteristics of 99 Cases of 2019-Novel Coronavirus (2019-nCoV) Pneumonia in Wuhan, China. (2020).
19. Channappanavar, R. & Perlman, S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin Immunopathol* **39**, 529-539 (2017).
20. Zhang, Y., *et al.* Analysis of serum cytokines in patients with severe acute respiratory syndrome. *Infect Immun* **72**, 4410-4415 (2004).
21. Sheng, W.H., *et al.* Clinical manifestations and inflammatory cytokine responses in patients with severe acute respiratory syndrome. *J Formos Med Assoc* **104**, 715-723 (2005).
22. Chien, J.Y., Hsueh, P.R., Cheng, W.C., Yu, C.J. & Yang, P.C. Temporal changes in cytokine/chemokine profiles and pulmonary involvement in severe acute respiratory syndrome. *Respirology* **11**, 715-722 (2006).
23. Cao, Y., *et al.* Nanopore sequencing: a rapid solution for infectious disease epidemics. *Sci China Life Sci* **62**, 1101-1103 (2019).
24. Kuba, K., *et al.* A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat Med* **11**, 875-879 (2005).
25. Zou, Z., *et al.* Angiotensin-converting enzyme 2 protects from lethal avian influenza A H5N1 infections. *Nat Commun* **5**, 3594 (2014).
26. de Jong MD1, S.C., Thanh TT, Hien VM, Smith GJ, Chau TN, Hoang DM, Chau NV, Khanh TH, Dong VC, Qui PT, Cam BV, Ha do Q, Guan Y, Peiris JS, Chinh NT, Hien TT, Farrar J.

- Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med* (2006).
27. Guo, J., *et al.* The Serum Profile of Hypercytokinemia Factors Identified in H7N9-Infected Patients can Predict Fatal Outcomes. *Sci Rep* **5**, 10942 (2015).
  28. Ferrari, S.M., *et al.* Novel therapies for thyroid autoimmune diseases: An update. *Best Pract Res Clin Endocrinol Metab*, 101366 (2019).
  29. Fallahi, P., *et al.* The aggregation between AITD with rheumatologic, or dermatologic, autoimmune diseases. *Best Pract Res Clin Endocrinol Metab*, 101372 (2019).
  30. Bi, Y.H., *et al.* Clinical and Immunological Characteristics of Human Infections With H5N6 Avian Influenza Virus. *Clin Infect Dis* **68**, 1100-1109 (2019).
  31. Pestka S, K.C., Walter MR. Interferons, interferon-like cytokines, and their receptors. *Immunol Rev* (2004).
  32. Shim JM, K.J., Tenson T, Min JY, Kainov DE. Influenza Virus Infection, Interferon Response, Viral Counter-Response, and Apoptosis. *Viruses* (2017).
  33. China National Health Commission Guidelines for Diagnosis and Treatment of 2019-nCoV Pneumonia. <http://www.gov.cn/zhengce/zhengceku/2020-01/28/5472673/files/0f96c10cc09d4d36a6f9a9f0b42d972b.pdf>. (2020).
  34. Mantovani, A., Barajon, I. & Garlanda, C. IL-1 and IL-1 regulatory pathways in cancer progression and therapy. *Immunol Rev* **281**, 57-61 (2018).
  35. Teng, M.W., *et al.* IL-12 and IL-23 cytokines: from discovery to targeted therapies for immune-mediated inflammatory diseases. *Nat Med* **21**, 719-729 (2015).
  36. Li, C.G., *et al.* IL-17 response mediates acute lung injury induced by the 2009 Pandemic Influenza A (H1N1) Virus. *Cell Res* **22**, 528-538 (2012).
  37. Wang, W., *et al.* Monoclonal antibody against CXCL-10/IP-10 ameliorates influenza A (H1N1) virus induced acute lung injury. *Cell Res* **23**, 577-580 (2013).
  38. Specific primers and probes for detection 2019 novel coronavirus. [http://ivdc.chinacdc.cn/kyjz/202001/t20200121\\_211337.html?from=timeline&isappinstall=0](http://ivdc.chinacdc.cn/kyjz/202001/t20200121_211337.html?from=timeline&isappinstall=0).
  39. Murray JF, M.M., Luce JM, Flick MR. An expanded definition of the adult respiratory distress syndrome. *Am Rev Respir Dis*. (1988).

## Acknowledgement

The authors would like to thank a scientist who insist to be anonymous. This work was supported by the National Natural Science Foundation of China (Grant 81788101) and the Chinese Academy of Medical Sciences Innovation Fund for Medical Sciences (Grant 2017-12M-1-009).

### **Author contributions**

C. Jiang conceived the project. C. Jiang, L. Liu, and W. J. Liu guided the study. Y. Liu, Y. Yang, F. Wang, J. Yuan, Z. Zhang, Z. Wang, J. Li, C. Shen, J. Li, L. Peng, W. Wu, M. Cao, L. Xing, Z. Xu, and L. Chen collected clinical samples from 2019-nCoV infected patients, H7N9-infected patients, bacterial pneumonia patients, and healthy subjects. C. Zhang, supervised by C. Zhou, measured cytokines/chemokines. F. Huang performed biostatistical analysis assisted by C. Zhang, Y. Qin, X. Li, D. Zhao, and S. Li. S. Tan made helpful assistant. C. Jiang, F. Huang, C. Zhang, and Y. Qin wrote the manuscript. All the authors have read and approved the manuscript.

### **Competing interests**

The authors declare no competing interests.

Table 1. Correlation between cytokine levels and viral Ct value of patients with laboratory-confirmed 2019-nCoV infection.

Cytokine	Ct value	
	Spearman	P value
M-CSF	-0.728	0.001
IL-10	-0.685	0.002
IFN- $\alpha$ 2	-0.653	0.003
IL-13	-0.636	0.005
IL-17	-0.608	0.007
IL-4	-0.593	0.009
IP-10	-0.568	0.014
IL-1 $\beta$	-0.521	0.026
IL-7	-0.514	0.029
IL-1ra	-0.514	0.029
G-CSF	-0.511	0.030
IL-12 (p40)	-0.511	0.030
IFN- $\gamma$	-0.508	0.031
IL-1 $\alpha$	-0.505	0.033
IL-2	-0.490	0.039
HGF	-0.470	0.049
PDGF-BB	0.677	0.002
MCP-3	-0.462	#
MIP-1 $\alpha$	-0.440	#
MCP-1 (MCAF)	-0.422	#
GM-CSF	-0.418	#
SCF	-0.372	-
IL-6	-0.367	-
MIG	-0.362	-
IL-8	-0.361	-
IL-18	-0.332	-
IL-3	-0.310	-
$\beta$ -NGF	-0.292	-
SCGF- $\beta$	-0.279	-
IL-12 (p70)	-0.278	-
FGF basic	-0.263	-
TRAIL	-0.247	-
TNF- $\alpha$	-0.208	-
IL-2R $\alpha$	-0.018	-
LIF	0.033	-
SDF-1 $\alpha$	0.124	-
TNF- $\beta$	0.128	-
CTACK	0.180	-

The viral titers were measured in 18 samples from 10 patients infected with 2019-nCoV. #: P value [0.05,0.1), - : P value >0.05

Fig. 1




Fig. 1 Murray Score highly correlates with plasma cytokine levels in patients with 2019-nCoV infections. The plasma levels of cytokines (IL-12, IFN- $\gamma$ , IL-2, HGF, IFN- $\alpha$ 2, IL-4, IL-17, IP-10, G-CSF, IL-10, IL-1ra, M-CSF, IL-1 $\alpha$ , IL-7 and PDGF-BB) were measured in a total of 25 blood samples from 12 patients with 2019-nCoV infections. Clinical indicators from the same day of blood sample collection were used to calculate Murray Score (an indicator of lung injury severity). Spearman rank correlation analysis (r) was used for linear correlation analysis.

Fig. 2

chinaXiv:202002.00018v1




Fig. 2 The ROC curve of plasma cytokine levels for patients with mild and severe 2019-nCoV-infections. The area under the receiver operating characteristic (ROC) curve (AUC) of the plasma cytokine levels (IL-12, IL-1ra, IP-10, PDGF-BB, IFN- $\alpha$ 2, IFN- $\gamma$ , M-CSF, IL-17, HGF, G-CSF, IL-2, IL-4, IL-10, IL-1 $\alpha$  and IL-7) was estimated in eight samples from four mildly infected 2019-nCoV- patients and 17 samples from eight severely infected patients. The P values of all AUC for plasma cytokine levels were less than 0.05.

Fig. 3




Fig. 3 Plasma cytokine levels in healthy controls, bacteria pneumonia patients, H7N9-infected patients and 2019-nCoV-infected patients The plasma levels of cytokines (M-CSF, IL-10, IFN-α2, IL-17, IL-4, IP-10, IL-7, IL-1ra, G-CSF, IL-12 (p40), IFN-γ, IL-1α, IL-2, HGF, and PDGF-BB) were measured in eight samples from eight healthy controls, eight samples from eight bacteria pneumonia patients, eight samples from eight H7N9-infected patients, eight samples from four 2019-nCoV-infected patients with mild illness (2019-nCoV-M) and 17 samples from eight patients with severe 2019-nCoV infection (2019-nCoV-S). Detailed information is shown in Extended Data Table 2.

\*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001.



Extended Data Table 1. Epidemiological and clinical features of subjects hospitalized with 2019-nCoV, H7N9 avian influenza virus and bacterial infections.

Characteristics	2019-nCoV	H7N9	Bacteria	Control
<b>Median age (range)</b>	62.5 (10-72)	56 (21-67)	41 (31.5-49)	28 (25-34)
<b>Age subgroups</b>				
0–15 years	1/12 (8.3%)	0/8 (0%)	1/8 (12.5%)	0/8 (0%)
16–59 years	4/12 (33.3%)	6/8 (75%)	5/8 (62.5%)	8/8 (100%)
≥60 years	7/12 (58.3%)	2/8 (25%)	2/8 (25%)	0/8 (0%)
<b>Male (%)</b>	8/12 (66.7%)	5/8 (62.5%)	5/8 (62.5%)	4/8 (50%)
<b>Initial symptoms</b>				
Fever	9/12 (75%)	8/8 (100%)	5/8 (62.5%)	0/8 (0%)
Cough	12/12 (100%)	7/8 (75%)	3/8 (37.5%)	0/8 (0%)
Headache	0/12 (0%)	3/8 (37.5%)	1/8 (12.5%)	0/8 (0%)
Myalgia	4/12 (33.3%)	4/8 (50%)	0/8 (0%)	0/8 (0%)
Chill	5/12 (41.7%)	5/8 (62.5%)	0/8 (0%)	0/8 (0%)
Nausea or vomiting	2/12 (16.7%)	0/8 (0%)	1/8 (12.5%)	0/8 (0%)
Diarrhea	3/12 (25%)	2/8 (25%)	0/8 (0%)	0/8 (0%)
<b>Co-existing chronic medical conditions</b>	6/12 (50%)	6/8 (50%)	5/8 (62.5%)	0/8 (0%)
Chronic heart disease	4/12 (33.3%)	4/8 (33.3%)	2/8 (25%)	0/8 (0%)
Chronic lung disease	1/12 (8.3%)	5/8 (41.7%)	1/8 (12.5%)	0/8 (0%)
Chronic renal disease	2/12 (16.7%)	2/8 (16.7%)	0/8 (0%)	0/8 (0%)
Chronic liver disease	1/12 (8.3%)	3/8 (25%)	0/8 (0%)	0/8 (0%)
Diabetes	2/12 (16.7%)	6/8 (50%)	0/8 (0%)	0/8 (0%)
Cancer	0/12 (0%)	4/8 (33.3%)	1/8 (12.5%)	0/8 (0%)
<b>Bacterial co-infections</b>	2/12 (16.7%)	7/8 (87.5%)	NA	0/8 (0%)
<b>Interval, median days (IQ+R)</b>				
Onset to admission	6.5 (5, 9.25)	6 (4.75, 7)	1 (0, 3.5)	NA
Onset to starting antiviral treatment	6 (5, 9.25)	5.5 (3.75, 6.25)	NA	NA
Onset to laboratory confirmation	7 (4.5, 14.5)	7 (5.75, 8.25)	6 (4.5, 8)	NA
<b>Complications</b>				
Pneumonia	12/12 (100%)	8/8 (100%)	8/8 (100%)	0/8 (0%)
ARDS	6/12 (50%)	7/8 (87.5%)	6/8 (75%)	0/8 (0%)
Severe ARDS	2/12 (16.7%)	5/8 (62.5%)	2/8 (25%)	0/8 (0%)
Respiratory failure	3/12 (25%)	7/8 (87.5%)	5/8 (62.5%)	0/8 (0%)
Hepatic insufficiency	2/12 (16.7%)	4/8 (50%)	4/8 (50%)	0/8 (0%)
Renal insufficiency	2/12 (16.7%)	2/8 (25%)	5/8 (62.5%)	0/8 (0%)
Cardiac failure	1/12 (8.3%)	3/8 (37.5%)	3/8 (37.5%)	0/8 (0%)
Shock	1/12 (8.3%)	1/8 (12.5%)	4/8 (50%)	0/8 (0%)
<b>Treatment</b>				
Received antivirals ≤ 2 days after illness onset	1/12 (8.3%)	1/8 (12.5%)	NA	NA
Received antivirals 3-5 days after illness onset	4/12 (33.3%)	3/8 (37.5%)	NA	NA
Received antivirals 6 days after illness onset	7/12 (58.3%)	4/8 (50%)	NA	NA
Corticosteroid	2/12 (16.7%)	8/8 (100%)	2/8 (25%)	NA
Mechanical ventilation	5/12 (41.7%)	7/8 (87.5%)	7/8 (87.5%)	NA

NA: Not applicable.

Extended Data Table 2. Cytokine comparison among healthy controls, bacteria-infected patients, H7N9-infected patients and 2019-nCoV-infected patients (disease severity classified)

Cytokine	Healthy						Bacteria					H7N9				2019 - nCov				
	Mean	SE	P <sup>a</sup> value	P <sup>b</sup> value	P <sup>c</sup> value	P <sup>d</sup> value	Mean	SE	P <sup>e</sup> value	P <sup>f</sup> value	P <sup>g</sup> value	Mean	SE	P <sup>h</sup> value	P <sup>i</sup> value	2019 - nCoV - M		2019 - nCoV - S		P <sup>j</sup> value
																Mean	SE	Mean	SE	
IL -1β	1.64	0.51	#	0.002	-	0.002	3.23	0.76	0.040	-	-	6.74	1.34	0.036	#	3.10	0.71	4.12	0.42	-
IL -1ra	156.74	7.40	0.001	0.001	<0.001	<0.001	5576.32	2956.33	-	0.002	-	11882.61	7235.11	0.016	-	283.15	35.42	1460.46	409.56	<0.001
IL -2	2.90	0.88	#	0.020	0.045	<0.001	6.89	2.21	-	-	0.041	12.16	3.64	-	-	5.46	0.38	9.71	0.88	0.002
IL -4	0.63	0.08	0.010	0.001	0.003	<0.001	1.43	0.25	-	-	#	2.03	0.41	-	-	1.27	0.12	1.92	0.15	0.007
IL -5	0.10	0.00	-	#	-	-	5.33	5.23	-	-	-	19.36	9.92	#	0.008	0.10	0.00	0.10	0.00	-
IL -6	4.87	1.68	0.006	-	-	0.004	201.55	109.57	-	0.002	-	199.98	130.85	#	-	3.26	0.60	22.79	6.93	<0.001
IL -7	10.01	0.67	-	0.020	0.002	<0.001	10.08	1.01	0.018	0.002	<0.001	17.84	2.67	-	#	18.05	1.42	25.26	1.81	0.012
IL -8	3.90	0.61	0.002	0.002	0.013	<0.001	74.98	32.66	-	0.021	-	102.02	53.69	0.006	-	6.53	0.55	18.80	2.84	0.001
IL -9	232.22	14.53	-	0.027	#	-	220.64	10.09	0.007	0.006	0.031	286.58	17.77	-	#	264.29	4.16	248.00	4.63	0.020
IL -10	0.89	0.66	-	0.009	0.011	<0.001	4.43	1.84	-	-	0.070	24.76	15.64	-	-	3.03	0.78	8.44	1.37	0.007
IL -12 (p70)	2.64	0.87	-	-	-	0.010	2.33	0.68	-	-	0.005	8.36	4.90	-	-	3.92	0.72	6.78	0.93	0.027
IL -13	2.73	0.66	-	0.026	0.008	<0.001	2.49	0.33	0.020	0.002	<0.001	8.49	2.44	-	-	5.98	0.80	8.01	0.52	0.041
IL -15	217.59	94.07	-	-	-	-	157.08	66.09	-	-	-	423.26	187.32	-	-	202.93	51.64	189.16	36.19	-
IL -17	7.12	1.66	0.045	0.015	-	0.003	11.59	1.55	-	-	-	14.69	2.46	0.036	-	9.06	0.52	14.24	1.02	<0.001
Eotaxin	30.68	5.20	-	-	-	#	41.14	10.93	-	-	-	56.77	13.86	-	-	37.23	2.56	44.67	4.74	-
FGF basic	30.80	1.82	-	0.020	-	<0.001	42.67	5.32	-	-	-	64.96	18.07	-	-	35.60	2.74	44.40	1.94	0.020
G-CSF	51.61	5.66	-	0.001	0.009	<0.001	367.40	232.15	-	-	-	321.14	114.59	<0.001	-	83.99	7.42	164.24	16.74	<0.001
GM-CSF	1.31	0.54	-	#	#	0.020	1.61	0.54	-	-	0.036	7.99	4.11	-	-	2.36	0.30	3.28	0.41	-
IFN-γ	5.21	1.12	0.010	0.001	0.006	<0.001	86.89	44.55	-	#	-	104.98	60.00	0.005	-	10.14	0.84	28.00	4.37	<0.001
IP-10	192.09	29.31	0.002	0.001	0.002	<0.001	3293.89	2007.72	0.046	-	-	22962.93	10848.70	<0.001	#	453.41	60.78	3386.70	631.17	<0.001
MCP-1 (MCAF)	8.78	1.91	#	0.024	0.036	<0.001	354.66	221.30	-	-	-	265.58	209.79	-	-	16.25	2.81	64.83	13.83	0.004
MIP-1α	1.32	0.09	0.008	0.001	0.005	<0.001	6.73	2.46	-	-	-	5.68	1.58	0.005	-	2.21	0.29	4.35	0.77	0.014
PDGF-BB	1972.26	305.97	#	-	-	0.010	1192.82	276.37	0.036	0.036	-	3675.14	1199.46	-	0.006	2157.76	239.56	1017.38	122.07	<0.001
MIP-1β	167.91	15.86	-	-	-	-	158.80	13.59	#	#	-	184.35	13.52	-	-	167.82	3.48	158.13	4.06	#
RANTES	4993.07	1385.58	0.046	0.027	0.021	-	7402.33	2024.74	-	-	#	14094.99	4181.73	-	0.012	6699.77	661.12	4677.52	494.74	0.014
TNF-α	53.19	4.47	0.036	0.006	0.036	<0.001	72.20	6.84	-	-	-	118.23	24.93	#	-	69.36	1.20	82.58	3.83	0.048
VEGF	105.72	27.11	-	0.046	-	-	114.11	24.96	0.045	-	-	265.98	76.46	0.036	0.027	108.66	9.21	130.88	10.77	#
IL -1a	5.05	2.75	-	0.002	0.005	<0.001	15.89	6.91	0.045	-	0.012	35.82	7.71	#	-	20.11	2.57	33.38	3.34	0.012
IL -2Rα	53.32	6.15	0.020	0.002	-	0.004	160.43	36.94	-	0.021	-	193.98	40.71	0.003	0.020	61.41	7.01	94.64	9.83	0.041
IL -3	0.23	0.08	-	#	#	0.003	0.33	0.14	-	-	0.006	1.38	0.82	-	-	0.44	0.10	0.74	0.10	0.044
IL -12 (p40)	71.51	11.89	0.001	0.002	0.003	<0.001	383.78	137.78	-	0.005	-	376.17	139.09	0.012	-	117.36	6.69	282.06	34.79	<0.001
IL -16	19.92	11.27	0.015	0.035	#	#	321.46	155.90	-	-	-	351.56	176.31	-	-	91.35	42.02	61.56	18.03	-
IL -18	28.90	7.51	0.002	0.002	0.005	<0.001	305.77	82.64	-	0.012	0.041	541.14	290.27	0.027	0.048	69.90	5.79	101.17	12.93	-
CTACK	310.34	76.56	0.005	0.003	0.021	0.005	1469.44	451.98	-	-	#	1154.40	252.81	#	0.041	558.71	64.27	653.19	85.09	-
GRO-α	937.91	160.22	-	-	-	-	610.87	39.31	-	0.016	-	786.16	99.14	-	-	730.27	25.44	652.08	18.89	#
HGF	137.42	15.75	0.001	0.002	-	<0.001	2375.54	1030.70	-	<0.001	0.023	2548.55	1476.82	0.001	-	166.35	9.81	872.79	278.63	<0.001
IFN-α2	8.30	1.84	-	0.004	0.027	<0.001	15.69	3.26	-	-	-	26.71	7.63	0.036	-	13.28	0.97	20.76	1.09	<0.001
LIF	0.10	0.00	0.011	0.027	0.027	0.001	29.48	16.43	-	-	-	64.48	27.74	-	-	18.25	8.25	23.87	5.79	-
MCP-3	1.78	0.89	-	0.004	#	<0.001	20.95	11.03	-	-	-	19.09	13.20	0.011	-	2.64	0.40	8.73	1.19	<0.001
M-CSF	14.30	1.47	0.003	0.001	0.003	<0.001	57.27	8.43	-	0.012	#	113.37	42.15	0.006	-	22.06	1.75	40.49	3.57	<0.001
MIF	1095.90	372.60	0.012	0.021	0.036	-	2466.29	664.87	-	-	0.017	4505.83	1729.28	-	#	1438.12	177.56	1153.54	117.92	-
MIG	63.20	12.69	0.010	0.001	0.016	<0.001	622.95	211.23	0.046	0.021	-	18787.15	13163.69	0.002	0.036	130.85	29.20	1088.53	315.64	0.003
b-NGF	0.64	0.24	#	0.006	0.013	<0.001	3.06	1.36	-	-	-	4.57	2.20	0.027	-	1.54	0.14	2.63	0.34	0.009
SCF	44.53	8.44	0.024	0.006	-	0.002	153.21	43.95	-	-	-	222.52	120.66	0.016	-	49.18	4.13	85.73	9.64	0.004
SCGF-β	48340.78	5710.60	0.002	0.001	#	0.006	140757.32	28033.76	-	0.009	-	176450.97	29405.62	0.002	0.012	64805.66	6401.71	99512.14	12073.24	-
SDF-1α	693.52	98.80	0.036	-	-	0.048	907.92	85.68	-	-	-	931.76	119.70	-	-	760.08	50.90	780.15	33.18	-
TNF-β	0.10	0.00	-	0.004	0.010	0.001	1.49	1.02	0.033	-	-	6.29	1.82	-	-	1.27	0.47	2.38	0.47	-
TRAIL	27.86	1.73	-	0.014	0.002	<0.001	32.92	3.90	-	-	0.027	58.34	20.27	-	-	37.62	1.50	43.87	2.16	#

P<sup>a</sup> value: Healthy vs bacteria patients
P<sup>b</sup> value: Healthy vs H7N9 patients
P<sup>c</sup> value: Healthy vs 2019-nCoV-M patients
P<sup>d</sup> value: Healthy vs 2019-nCoV-S patients
P<sup>e</sup> value: Bacteria patients vs H7N9 patients

P<sup>f</sup> value: Bacteria patients vs 2019-nCoV-M patients
P<sup>g</sup> value: Bacteria patients vs 2019-nCoV-S patients
P<sup>h</sup> value: H7N9 patients vs 2019-nCoV-M patients
P<sup>i</sup> value: H7N9 patients vs 2019-nCoV-S patients
P<sup>j</sup> value : 2019-nCoV-M patients vs 2019-nCoV-S patients

# : P value [0.05,0.1), - : P value >0.05

8 samples from 8 healthy control, 8 samples from 8 bacteria-infected patients, 8 samples from 8 H7N9-infected patients, 8 samples from 4 mild 2019-nCoV-infected patients (2019-nCoV-M) and 17 samples from 8 severe 2019-nCoV-infected patients (2019-nCoV-S).

Extended Data Table 3. Cytokine comparison among healthy controls, bacteria-infected patients, H7N9-infected patients and 2019-nCoV-infected patients (Day0-7, Day8-14, Day15-)

Cytokine	Healthy			Bacteria			H7N9			2019-nCoV								
	P <sup>a</sup> value	P <sup>b</sup> value	P <sup>c</sup> value	P <sup>d</sup> value	P <sup>e</sup> value	P <sup>f</sup> value	P <sup>g</sup> value	P <sup>h</sup> value	P <sup>i</sup> value	Day 0-7		Day 8-14		Day 15-		P <sup>j</sup> value	P <sup>k</sup> value	P <sup>l</sup> value
										Mean	SE	Mean	SE	Mean	SE			
IL-1β	0.037	0.020	0.019	-	-	-	-	0.046	#	4.08	1.02	3.63	0.48	3.35	0.49	-	-	-
IL-1ra	0.001	<0.001	0.003	#	0.011	-	-	-	-	1478.23	917.46	889.43	308.43	867.02	259.33	-	-	-
IL-2	0.027	0.002	0.005	-	-	-	-	-	-	6.44	0.84	8.91	1.12	7.74	0.68	-	#	-
IL-4	0.004	0.001	0.003	-	-	-	-	-	-	1.32	0.15	1.90	0.18	1.47	0.10	0.043	-	-
IL-5	-	-	-	-	-	-	#	0.020	-	0.10	0.00	0.10	0.00	0.10	0.00	-	-	-
IL-6	-	#	#	0.028	0.017	-	-	-	-	9.98	3.54	10.07	2.22	34.62	22.56	-	#	-
IL-7	0.003	0.001	0.003	0.003	<0.001	0.003	-	#	-	17.66	1.46	25.54	2.10	19.65	1.02	0.017	-	-
IL-8	0.008	0.001	0.003	#	-	-	0.049	-	-	9.87	2.79	15.85	3.24	16.70	6.49	-	-	-
IL-9	-	-	-	0.049	0.030	0.019	-	-	-	254.65	6.41	248.48	5.69	263.36	5.75	-	-	-
IL-10	0.006	0.001	0.009	-	-	-	-	-	-	6.45	2.79	7.05	1.36	4.33	1.17	-	-	-
IL-12 (p70)	-	0.014	0.040	-	0.012	0.012	-	-	-	3.62	0.34	6.74	1.12	4.78	0.53	-	0.034	-
IL-13	0.011	0.001	0.008	0.002	<0.001	0.003	-	-	-	6.61	0.90	7.65	0.67	6.86	1.06	-	-	-
IL-15	-	-	-	-	-	-	-	-	-	104.65	43.18	218.73	39.35	190.10	66.90	#	-	-
IL-17	#	0.008	0.040	-	-	-	-	-	-	10.26	0.99	13.49	1.33	11.56	0.67	-	-	-
Eotaxin	#	-	-	-	-	-	-	-	-	44.86	6.72	38.29	4.69	45.29	8.15	-	-	-
FGF basic	-	0.003	0.008	-	-	-	-	-	-	37.23	3.52	43.14	2.51	39.63	3.85	-	-	-
G-CSF	0.005	0.001	0.003	-	-	-	#	0.036	-	107.31	14.62	148.44	23.14	136.99	26.02	-	-	-
GM-CSF	#	0.038	#	-	#	#	-	-	-	2.43	0.36	3.15	0.48	2.37	0.38	-	-	-
IFN-γ	0.005	0.001	0.003	-	-	-	-	-	-	19.65	8.00	22.36	4.59	22.20	7.14	-	-	-
IP-10	0.002	<0.001	0.003	-	-	-	#	0.007	#	2680.72	1294.44	1946.64	529.30	2944.85	1304.24	-	-	-
MCP-1 (MCAF)	0.028	0.001	0.008	-	-	-	-	-	-	34.48	11.44	41.98	10.99	74.12	40.87	-	-	-
MIP-1α	0.009	<0.001	0.004	-	-	-	0.049	-	-	2.54	0.35	4.22	1.00	3.63	0.98	-	-	-
PDGF-BB	-	#	-	-	-	-	-	0.025	-	1651.44	426.27	1241.39	170.50	1370.23	266.40	-	-	-
MIP-1β	-	-	-	-	-	-	-	-	-	160.45	3.18	159.07	5.19	169.04	5.34	-	-	-
RANTES	-	-	-	-	-	-	#	0.043	-	5683.14	925.79	4763.56	456.24	6404.49	1272.99	-	-	-
TNF-α	#	0.002	0.003	-	-	#	-	-	-	70.85	4.36	81.14	4.78	76.86	4.51	-	0.042	-
VEGF	-	-	-	-	-	-	#	0.030	#	112.19	12.95	126.58	12.13	113.28	9.97	-	-	-
IL-1a	0.006	<0.001	0.005	-	0.017	#	-	-	-	21.46	3.21	32.58	3.78	22.07	3.02	#	-	-
IL-2Rα	-	0.014	#	#	-	-	0.008	0.017	#	68.48	11.31	91.07	12.53	89.23	10.26	-	-	-
IL-3	-	0.003	#	-	0.014	0.039	-	-	-	0.38	0.05	0.77	0.11	0.45	0.05	0.017	-	-
IL-12 (p40)	0.004	<0.001	0.003	0.049	-	-	-	-	-	156.36	29.38	263.82	46.56	207.24	51.43	-	#	-
IL-16	0.049	-	0.035	-	-	-	-	-	-	56.54	15.36	48.52	18.26	154.14	65.84	-	-	-
IL-18	0.004	0.001	0.008	#	0.017	-	-	0.025	-	111.64	28.12	79.63	8.04	95.26	12.86	-	-	-
CTACK	#	0.006	0.019	-	-	-	#	#	-	496.00	83.70	641.32	88.69	728.05	174.99	-	-	-
GRO-α	-	-	-	-	-	0.040	-	-	-	669.95	20.46	657.17	23.28	723.07	51.58	-	-	-
HGF	0.037	0.006	0.013	0.005	0.025	0.019	0.028	#	#	305.64	121.51	875.59	354.58	518.05	322.31	-	-	-
IFN-α2	0.015	0.001	0.012	-	-	-	-	-	-	16.63	2.47	18.58	1.37	17.93	1.02	-	-	-
LIF	0.047	0.001	0.018	-	-	-	-	-	-	12.58	7.97	28.11	6.11	5.77	4.54	#	-	-
MCP-3	0.036	0.003	0.012	-	-	-	-	-	-	4.43	1.35	7.29	1.35	6.81	2.79	-	#	-
M-CSF	0.001	0.001	0.003	-	0.017	#	-	#	-	39.97	8.50	32.22	3.53	32.31	3.97	-	-	-
MIF	#	-	-	-	0.020	-	-	#	-	1295.66	173.44	1116.72	113.22	1511.60	325.36	-	-	-
MIG	0.011	0.001	0.005	-	-	-	0.021	0.020	#	469.21	258.66	939.15	390.93	749.79	489.35	-	-	-
b-NGF	0.017	0.002	0.005	-	-	-	-	-	-	1.49	0.18	2.55	0.42	2.04	0.25	0.047	0.015	-
SCF	-	0.007	0.019	-	-	-	-	-	-	58.44	8.34	78.61	12.45	78.51	16.59	-	-	-
SCGF-β	0.037	0.017	#	-	#	-	0.028	0.005	0.040	84399.44	14867.23	93153.33	14483.77	78223.91	18421.01	-	-	-
SDF-1α	-	-	#	-	-	-	-	-	-	761.70	52.72	750.62	33.61	846.56	78.83	-	-	-
TNF-β	-	0.001	0.001	-	#	#	#	-	-	0.91	0.57	2.35	0.51	1.62	0.52	#	#	-
TRAIL	0.002	0.001	0.003	#	#	-	-	-	-	43.32	3.81	39.70	1.77	45.13	4.31	-	-	-

P<sup>a</sup> value: Healthy vs 2019-nCoV patients (Day0-7)

P<sup>b</sup> value: Healthy vs 2019-nCoV patients (Day8-14)

P<sup>c</sup> value: Healthy vs 2019-nCoV patients (Day15-)

# : P value [0.05,0.1), - : P value >0.05

P<sup>d</sup> value: Bacteria patients vs 2019-nCoV patients (Day0-7)

P<sup>e</sup> value: Bacteria patients vs 2019-nCoV patients (Day8-14)

P<sup>f</sup> value: Bacteria patients vs 2019-nCoV patients (Day15-)

P<sup>g</sup> value: H7N9 patients vs 2019-nCoV patients (Day0-7)

P<sup>h</sup> value: H7N9 patients vs 2019-nCoV patients (Day8-14)

P<sup>i</sup> value: H7N9 patients vs 2019-nCoV patients (Day15-)

P<sup>j</sup> value : 2019-nCoV patients (Day0-7) vs 2019-nCoV patients (Day8-14)

P<sup>k</sup> value : 2019-nCoV patients (Day0-7) vs 2019-nCoV patients (Day15-)

P<sup>l</sup> value : 2019-nCoV patients (Day8-14) vs 2019-nCoV patients (Day15-)

8 samples from 8 healthy control, 8 samples from 8 bacteria-infected patients, 8 samples from 8 H7N9-infected patients, 7 samples from 7 2019-nCoV-infected patients (Day0-7), 13 samples from 9 2019-nCoV-infected patients (Day8-14) and 5 samples from 5 2019-nCoV-infected patients (Day15-).

Extended Data Table 4. Detection of 2019-nCoV at respiratory and non-respiratory sites.

		2019-nCoV		
		Total	Severe	Mild
<b>Nasopharynx</b>				
Detectable RNA (n/N)	Throat	8/11	5/7	3/4
	Sputum	7/10	6/6	1/4
Ct Values (Median; range)	Throat	30 (23-36)	29.2 (25-36)	30 (23-35)
	Sputum	29.1 (20-34.7)	28.3 (20-33.4)	34.3
<b>BLAF</b>				
Detectable RNA (n/N)		3/5	3/4	0/1
Ct Values (Median; range)		24 (19-26)	24 (19-26)	U
<b>Plasma</b>				
Detectable RNA (n/N)		0/12	0/8	0/4
Ct Values (Median; range)		U	U	U
<b>Rectum</b>				
Detectable RNA (n/N)		2/7	1/5	1/2
Ct Values (Median;range)		27.5 (27-28)	28	27

U: Undetected.

Throat swabs were obtained after 4–18 d (median 10) of illness.

Sputum were obtained after 4–17 d (median 9.5) of illness.

Plasma samples were obtained after 7–10 d (median 9) of illness.

Rectal swabs were obtained after 4–16.5 d (median 9.5) of illness.